

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number
WO 03/088978 A1

(51) International Patent Classification⁷: A61K 31/7076,
31/138, 31/44, A61P 9/06

(21) International Application Number: PCT/US03/12043

(22) International Filing Date: 18 April 2003 (18.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/373,766 18 April 2002 (18.04.2002) US

(71) Applicant: CV THERAPEUTICS, INC. [US/US]; 3172
Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors: BELARDINELLI, Luiz; 704 University
Drive, Menlo Park, CA 94025 (US). DHALLA, Arvin-
der; 131 College Avenue, Mountain View, CA 94040
(US).

(74) Agents: CLARKE, Pauline et al.; CV Therapeutics, Inc.,
3172 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF TREATING ARRHYTHMIAS COMPRISING ADMINISTRATION OF AN A1 ADENOSINE AGO-
NIST WITH A BETA BLOCKER, CALCIUM CHANNEL BLOCKER OR A CARDIAC GLYCOSIDE



(57) Abstract: ABSTRACT Methods are provided for treating arrhythmia in a manner that minimizes undesirable side effects, comprising administration of a therapeutically effective minimal dose of an A1 adenosine receptor agonist with a therapeutically effective minimal dose of a beta blocker, calcium channel blocker, or a cardiac glycoside.

METHOD OF TREATING ARRHYTHMIAS COMPRISING ADMINISTRATION OF AN A₁ ADENOSINE AGONIST WITH A BETA BLOCKER, CALCIUM CHANNEL BLOCKER OR A CARDIAC GLYCOSIDE

Priority is claimed to U.S. Provisional Patent Application Serial No. 06/373,766, filed April 19, 2002, the complete disclosure of which is hereby incorporated by reference.

5 Field of the Invention

This invention relates to a method of treating arrhythmias and heart failure in a manner that minimizes undesirable side effects, comprising administration of a low dose of an adenosine A₁ receptor agonist in conjunction with a low dose of a beta blocker, or a calcium channel blocker, or a cardiac glycoside.

10

Background Information

Arrhythmias are abnormal heart rhythms that occur either in the atria or the ventricles. Arrhythmias arising in the atria are called atrial arrhythmias, and these disorders include atrial fibrillation, atrial flutter, and paroxysmal atrial tachycardia (PSVT). Arrhythmias arising in the ventricles, known as ventricular arrhythmias, are a group of disorders having diverse etiologies, including idiopathic ventricular tachycardia, ventricular fibrillation, and Torsade de Pointes (TdP). Arrhythmias can range from incidental, asymptomatic clinical findings to life-threatening abnormalities, and account for a significant percentage of the causes of death in humans. Thus, it is desirable to develop methods of mitigating the effects of arrhythmias.

20 A variety of anti-arrhythmic drug therapies are presently available, and are classified as follows. Class I anti-arrhythmics, comprising sodium channel blockers; Class II, comprising beta-blockers; Class III, comprising drugs that prolong action potential (usually by blocking potassium channels); and Class IV, comprising calcium channels blockers. Cardiac glycosides, for example digitalis, are also used as drugs for the treatment of arrhythmia, but they have a delayed onset of action (about 30 minutes) and their peak effects are not observed for ≥ 3 to 4 hours after administration. Additionally, digitalis is toxic at doses close to the therapeutic dose, which limits the utility of the compound.

30 In fact all of the above classes have significant limitations. For example, beta-blockers, such as propranolol and esmolol, and calcium-channel blockers, for example verapamil, bepridil, and diltiazem, can cause hypotension, potentially have negative inotropic effects, and may also precipitate new arrhythmias, including TdP.

35 Adenosine, which is widely found in nature, is another compound that has anti-arrhythmic activities, by virtue of its ability, at certain dose levels, to slow the conduction in the atrioventricular node. The anti-arrhythmic effects of adenosine are due exclusively to its interaction with the adenosine A₁ receptor subtype. However, although adenosine is highly

effective in ameliorating arrhythmia, it also binds contemporaneously to other adenosine receptor subtypes (A_{2A} , A_{2B} , and A_3), which results in undesirable side effects, such as vasodilation, changes in the heart rate, mast cell degradation, etc. Adenosine also has a short half-life (~10 sec), making it ineffective in treating conditions that require prolonged action.

5 Compounds that are selective agonists for adenosine A_1 receptors are known. For example, a new class of agonists that bind to adenosine A_1 receptors and that are useful in treating arrhythmias are disclosed in U.S. Patent No. 5,789,416, and in U.S. Patent Application, Serial No. 10/194,335, the entire disclosures of which are hereby incorporated by reference. These compounds have a high specificity for the adenosine A_1 receptor subtype, but like all
10 therapeutic compounds, can potentially cause side effects.

 Antiarrhythmic agents in general have a narrow margin between the dose required to produce the desired antiarrhythmic effect and the dose that produces an adverse effect. It would therefore be desirable to find a method of treating arrhythmia that is effective at low doses (or minimal doses) of the active agent, thus decreasing the likelihood of adverse effects. We have
15 discovered that low doses of adenosine A_1 receptor agonists, preferably partial agonists, and more preferably selective adenosine A_1 receptor agonists, can be used in combination with low doses of beta blocker, calcium channel blockers, or cardiac glycosides, to provide an effective treatment for arrhythmia that minimizes the side effects of beta blockers, calcium channel blockers, cardiac glycosides, and A_1 -adenosine receptor agonists that may potentially occur
20 when taken individually. It has also been observed that at low doses, the combination of these agents act in a synergistic manner, thus reducing even further the chance of side effects. It has also been observed that the combination of an A_1 adenosine receptor antagonist with a beta blocker can be used in the treatment of heart failure, including ischemic heart disease, congestive heart failure, heart failure syndrome, hypertension, and the like.

25 Accordingly, a novel and effective method of treating arrhythmias is provided that restores sinus rhythm without slowing the sinus rate and is virtually free of undesirable side effects, such as changes in mean arterial pressure, blood pressure, increased heart rate, TdP, or other adverse effects.


30

SUMMARY OF THE INVENTION

 It is an object of this invention to provide an effective method of treating arrhythmias in a mammal while minimizing undesirable side effects. Accordingly, in a first aspect, the invention relates to a method of treating arrhythmias in a mammal, comprising administration of a
35 therapeutically effective minimal dose of an A_1 adenosine receptor agonist in conjunction with a

therapeutically effective minimal dose of a beta blocker, calcium channel blocker, or a cardiac glycoside to a mammal in need thereof.

In one embodiment, an A₁-adenosine receptor agonist useful for this invention is a compound of Formula I:



Formula I


wherein:

R₁ is an optionally substituted heterocyclic group, preferably monocyclic.

The effective dose is preferably in the range of 0.0001-0.05 mg/kg, more preferably 0.0005-0.02 mg/kg.

In a preferred embodiment, R¹ is 3-tetrahydrofuranyl, 3-tetrahydrothiofuranyl, 4-pyranyl, or 4-thiopyranyl. The most preferred compound of Formula I is 6-(3-(R)-N-aminotetrahydrofuranyl)purine riboside (hereinafter referred to as CVT-510).

In another embodiment, an A₁-adenosine receptor agonist useful for this invention is a compound of Formula II:



Formula II

wherein:

R¹ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

R² is hydrogen, halo, trifluoromethyl, optionally substituted acyl, or cyano;

R³ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl;

5 optionally substituted heteroaryl, or optionally substituted heterocyclyl,

R⁴ and R⁵ are independently hydrogen or optionally substituted acyl; and

X and Y are independently a covalent bond or optionally substituted alkylene.

The effective dose is preferably in the range of 0.1 to 200 mg/kg, more preferably 0.5 to 50 mg/kg.

10 The most preferred compound of Formula II is one in which R¹ is 2-hydroxycyclopentyl, X and Y are covalent bonds, R², R³, and R⁴ are hydrogen, and R⁵ is 2-fluorophenyl, most preferably 2-{6-[(1R,2R)-2-hydroxycyclopentyl]amino}purin-9-yl}(4S,5S,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol, hereinafter referred to as CVT-3619.

Preferred beta blockers include atenolol, esmolol, sotalol, and propranolol. More preferred is esmolol. The preferred effective dose is in the range of 0.01 to 100 mg/kg, more preferred in the range of 0.1 to 10 mg/kg.

Preferred calcium channel blockers include amlodipine, bepridil, diltiazem, felodipine, isradipine, nicardipine, nifedipine, nimodipine and verapamil. The preferred effective dose is in the range of 0.01 to 50 mg/kg, more preferably in the range of 0.1 to 10 mg/kg.

20 Preferred cardiac glycosides include digoxin and digitoxin.

One preferred embodiment of the invention is a method of treating arrhythmias in a mammal comprising administering a therapeutically effective minimal dose of CVT-510 in conjunction with a therapeutically effective minimal dose of a beta blocker, preferably esmolol, atenolol, sotalol or propranolol, more preferably esmolol, to a mammal in need thereof.

25 Another preferred embodiment of the invention is a method of treating arrhythmias in a mammal comprising administering a therapeutically effective minimal dose of CVT-3619, and a therapeutically effective minimal dose of a beta blocker, preferably esmolol, atenolol, sotalol or propranolol, more preferably esmolol, to a mammal in need thereof.

Another preferred embodiment of the invention is a method of treating arrhythmias in a mammal comprising administering a therapeutically effective minimal dose of CVT-510, or a therapeutically effective minimal dose of CVT-3619, in conjunction with a therapeutically effective minimal dose of a calcium channel blocker, preferably verapamil, to a mammal in need thereof.

30 A third preferred embodiment of the invention is a method of treating arrhythmias in a mammal comprising administering a therapeutically effective minimal dose of CVT-510, or a

therapeutically effective minimal dose of CVT-3619, in conjunction with a therapeutically effective minimal dose of a cardiac glycoside, preferably digoxin, to a mammal in need thereof.

In another aspect, the invention relates to a pharmaceutical composition useful for treating arrhythmias in a mammal, comprising a therapeutically effective minimal dose of an A₁ adenosine receptor agonist and a therapeutically effective minimal dose of a beta blocker, and at least one pharmaceutically acceptable excipient.

One preferred embodiment of the invention is a pharmaceutical composition for treating arrhythmias in a mammal, comprising a therapeutically effective minimal dose of a compound of Formula I, more preferably CVT-510, and a therapeutically effective minimal dose of a beta blocker, preferably esmolol, atenolol, sotalol or propranolol, more preferably esmolol, to a mammal in need thereof. The Formula I dose is preferably in the range of 0.0001-0.05 mg/kg, more preferably 0.0005-0.02 mg/kg, and the beta blocker dose is in the in the range of 0.01 to 100 mg/kg, more preferably in the range of 0.1 to 10 mg/kg.

Another preferred embodiment of the invention is a pharmaceutical composition for treating arrhythmias in a mammal, comprising a therapeutically effective minimal dose of a compound of Formula II, more preferably CVT-3619, and a therapeutically effective minimal dose of a beta blocker, preferably esmolol, atenolol, sotalol or propranolol, more preferably esmolol, to a mammal in need thereof. The dose is preferably in the range of 0.1 to 200 mg/kg, more preferably 0.5 to 25 mg/kg, and the beta blocker dose is in the in the range of 0.01 to 100 mg/kg, more preferably in the range of 0.1 to 10 mg/kg.

In another aspect, the invention relates to a method of treating heart failure in a mammal, comprising administration of a therapeutically effective minimal dose of an A₁ adenosine receptor agonist in conjunction with a therapeutically effective minimal dose of a beta blocker to a mammal in need thereof.

DESCRIPTION OF FIGURES.

Figure 1. Comparison of the effect of CVT-3619 alone and CVT-3619 in combination with propranolol on heart rate.

Figure 2. Comparison of the effect of CVT-3619 alone and CVT-3619 in combination with propranolol on heart rate.

Figure 3. Comparison of the effect of 20µg/kg of CVT-510, 10mg/kg of esmolol, and a combination of 20µg/kg of CVT-510 and 10mg/kg of esmolol on heart rate.

Figure 4. Comparison of the effect of 20µg/kg of CVT-510, 3mg/kg of esmolol, and a combination of 20µg/kg of CVT-510 and 3mg/kg of esmolol on heart rate.

Figure 5. Comparison of the effect of 10 μ g/kg, 20 μ g/Kg and 30 μ g/kg doses of CVT-510, 1 mg/kg and 3mg/kg of esmolol, and a combination of 20 μ g/kg of CVT-510 and 1 and 3mg/kg of esmolol on duration of Bradycardia.

Figure 6. Comparison of plasma levels of CVT-510 alone and a combination of CVT-510 and metoprolol.

Figure 7. Dose response curve for metoprolol in the absence and presence of CVT-510.

Figure 8. This figure represents the data shown in figure 7.

Figure 9. Effect of CVT-510 (0.5 μ g/kg) and metoprolol (0.1 mg/kg), which were given as an iv bolus, on PR interval

Figure 10. Effect of CVT-510 (0.5 μ g/kg) and esmolol on PR interval

ABBREVIATIONS

BPM: beats per minute

HR: Heart rate

SH: Stimulus to His (length of time for conduction of current through AV node)

PSVT Paroxysmal Atrial Tachycardia

DEFINITIONS AND GENERAL PARAMETERS

As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

The term "alkyl" refers to a monoradical branched or unbranched saturated hydrocarbon chain having from 1 to 20 carbon atoms. This term is exemplified by groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, t-butyl, n-hexyl, n-decyl, tetradecyl, and the like.

The term "substituted alkyl" refers to:

- 1) an alkyl group as defined above, having from 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. Unless otherwise constrained by the definition, all substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2; or

- 2) an alkyl group as defined above that is interrupted by 1-5 atoms or groups independently chosen from oxygen, sulfur and $\text{-NR}_a\text{-}$, where R_a is chosen from hydrogen, alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, aryl, heteroaryl and heterocyclyl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF_3 , amino, substituted amino, cyano, or $\text{-S(O)}_n\text{R}$, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2; or
- 3) an alkyl group as defined above that has both from 1 to 5 substituents as defined above and is also interrupted by 1-5 atoms or groups as defined above.

The term "lower alkyl" refers to a monoradical branched or unbranched saturated hydrocarbon chain having from 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, t-butyl, n-hexyl, and the like.

The term "substituted lower alkyl" refers to lower alkyl as defined above having 1 to 5 substituents, preferably 1 to 3 substituents, as defined for substituted alkyl, or a lower alkyl group as defined above that is interrupted by 1-5 atoms as defined for substituted alkyl, or a lower alkyl group as defined above that has both from 1 to 5 substituents as defined above and is also interrupted by 1-5 atoms as defined above.

The term "alkylene" refers to a diradical of a branched or unbranched saturated hydrocarbon chain, preferably having from 1 to 20 carbon atoms, preferably 1-10 carbon atoms, more preferably 1-6 carbon atoms. This term is exemplified by groups such as methylene ($\text{-CH}_2\text{-}$), ethylene ($\text{-CH}_2\text{CH}_2\text{-}$), the propylene isomers (e.g., $\text{-CH}_2\text{CH}_2\text{CH}_2\text{-}$ and $\text{-CH(CH}_3\text{)CH}_2\text{-}$) and the like.

The term "lower alkylene" refers to a diradical of a branched or unbranched saturated hydrocarbon chain, preferably having from 1 to 6 carbon atoms.

The term "substituted alkylene" refers to:

- (1) an alkylene group as defined above having from 1 to 5 substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl , -SO-aryl , -SO-heteroaryl , $\text{-SO}_2\text{-alkyl}$, $\text{SO}_2\text{-aryl}$ and $\text{-SO}_2\text{-heteroaryl}$. Unless otherwise constrained by the definition, all substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF_3 , amino, substituted amino, cyano, or $\text{-S(O)}_n\text{R}$, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2; or
- (2) an alkylene group as defined above that is interrupted by 1-5 atoms or groups independently chosen from oxygen, sulfur and $\text{NR}_a\text{-}$, where R_a is chosen from hydrogen,

optionally substituted alkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl and heterocyclyl, or groups selected from carbonyl, carboxyester, carboxyamide and sulfonyl; or

- (3) an alkylene group as defined above that has both from 1 to 5 substituents as defined above and is also interrupted by 1-20 atoms as defined above. Examples of substituted alkylenes are chloromethylene (-CH(Cl)-), aminoethylene (-CH(NH₂)CH₂-), methylaminoethylene (-CH(NHMe)CH₂-), 2-carboxypropylene isomers (-CH₂CH(CO₂H)CH₂-), ethoxyethyl (-CH₂CH₂O-CH₂CH₂-), ethylmethylaminoethyl (-CH₂CH₂N(CH₃)CH₂CH₂-), 1-ethoxy-2-(2-ethoxy-ethoxy)ethane (-CH₂CH₂O-CH₂CH₂-OCH₂CH₂-OCH₂CH₂-), and the like.

The term "aralkyl" refers to an aryl group covalently linked to an alkylene group, where aryl and alkylene are defined herein. "Optionally substituted aralkyl" refers to an optionally substituted aryl group covalently linked to an optionally substituted alkylene group. Such aralkyl groups are exemplified by benzyl, phenylethyl, 3-(4-methoxyphenyl)propyl, and the like.

The term "alkoxy" refers to the group R-O-, where R is optionally substituted alkyl or optionally substituted cycloalkyl, or R is a group -Y-Z, in which Y is optionally substituted alkylene and Z is; optionally substituted alkenyl, optionally substituted alkynyl; or optionally substituted cycloalkenyl, where alkyl, alkenyl, alkynyl, cycloalkyl and cycloalkenyl are as defined herein. Preferred alkoxy groups are alkyl-O- and include, by way of example, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, 1,2-dimethylbutoxy, and the like.

The term "alkylthio" refers to the group R-S-, where R is as defined for alkoxy.

The term "alkenyl" refers to a monoradical of a branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 20 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having 1-6, preferably 1, double bond (vinyl). Preferred alkenyl groups include ethenyl or vinyl (-CH=CH₂), 1-propylene or allyl (-CH₂CH=CH₂), isopropylene (-C(CH₃)=CH₂), bicyclo[2.2.1]heptene, and the like. In the event that alkenyl is attached to nitrogen, the double bond cannot be alpha to the nitrogen.

The term "lower alkenyl" refers to alkenyl as defined above having from 2 to 6 carbon atoms.

The term "substituted alkenyl" refers to an alkenyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycloxy,

hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

5 The term "alkynyl" refers to a monoradical of an unsaturated hydrocarbon, preferably having from 2 to 20 carbon atoms, more preferably 2 to 10 carbon atoms and even more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-6 sites of acetylene (triple bond) unsaturation. Preferred alkynyl groups include ethynyl, (-C≡CH), propargyl (or propynyl, -C≡CCH₃), and the like. In the event that alkynyl is attached to nitrogen, the triple
10 bond cannot be alpha to the nitrogen.

 The term "substituted alkynyl" refers to an alkynyl group as defined above having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl,
15 carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or
20 heteroaryl and n is 0, 1 or 2.

 The term "aminocarbonyl" refers to the group -C(O)NRR where each R is independently hydrogen, alkyl, aryl, heteroaryl, heterocyclyl or where both R groups are joined to form a heterocyclic group (e.g., morpholino). All substituents may be optionally further substituted by
25 alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

 The term "acylamino" refers to the group -NRC(O)R where each R is independently hydrogen, alkyl, aryl, heteroaryl, or heterocyclyl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in
which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

30 The term "acyloxy" refers to the groups -O(O)C-alkyl, -O(O)C-cycloalkyl, -O(O)C-aryl, -O(O)C-heteroaryl, and -O(O)C-heterocyclyl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in
which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

The term "aryl" refers to an aromatic carbocyclic group of 6 to 20 carbon atoms having a single ring (e.g., phenyl) or multiple rings (e.g., biphenyl), or multiple condensed (fused) rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the like.

Unless otherwise constrained by the definition for the aryl substituent, such aryl groups
5 can optionally be substituted with from 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy,
10 heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

The term "aryloxy" refers to the group aryl-O- wherein the aryl group is as defined
15 above, and includes optionally substituted aryl groups as also defined above. The term "arylthio" refers to the group R-S-, where R is as defined for aryl.

The term "amino" refers to the group -NH₂.

The term "substituted amino" refers to the group -NRR where each R is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, carboxyalkyl (for example,
20 benzyloxycarbonyl), aryl, heteroaryl and heterocyclyl provided that both R groups are not hydrogen, or a group -Y-Z, in which Y is optionally substituted alkylene and Z is alkenyl, cycloalkenyl, or alkynyl,. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

25 The term "carboxyalkyl" refers to the groups -C(O)O-alkyl, -C(O)O-cycloalkyl, where alkyl and cycloalkyl, are as defined herein, and may be optionally further substituted by alkyl, alkenyl, alkynyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

The term "cycloalkyl" refers to cyclic alkyl groups of from 3 to 20 carbon atoms having a
30 single cyclic ring or multiple condensed rings. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and bicyclo[2.2.1]heptane, or cyclic alkyl groups to which is fused an aryl group, for example indan, and the like.

The term "substituted cycloalkyl" refers to cycloalkyl groups having from 1 to 5
35 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkyl,

alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2.

The term "halogen" or "halo" refers to fluoro, bromo, chloro, and iodo.

10 The term "acyl" denotes a group -C(O)R, in which R is hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heterocyclyl, optionally substituted aryl, and optionally substituted heteroaryl.

The term "heteroaryl" refers to an aromatic group (i.e., unsaturated) comprising 1 to 15 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within at least one ring.

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 5 substituents, preferably 1 to 3 substituents selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indoliziny, benzothiazole, or benzothienyl). Examples of nitrogen heterocycles and heteroaryls include, but are not limited to, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, and the like as well as N-alkoxy-nitrogen containing heteroaryl compounds.

The term "heteroaryloxy" refers to the group heteroaryl-O-.

35 The term "heterocyclyl" refers to a monoradical saturated or partially unsaturated group

having a single ring or multiple condensed rings, having from 1 to 40 carbon atoms and from 1 to 10 hetero atoms, preferably 1 to 4 heteroatoms, selected from nitrogen, sulfur, phosphorus, and/or oxygen within the ring.

Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5, and preferably 1 to 3 substituents, selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, amino, aminocarbonyl, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, thiocarbonyl, carboxy, carboxyalkyl, arylthio, heteroarylthio, heterocyclylthio, thiol, alkylthio, aryl, aryloxy, heteroaryl, aminosulfonyl, aminocarbonylamino, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, SO₂-aryl and -SO₂-heteroaryl. All substituents may be optionally further substituted by alkyl, alkoxy, halogen, CF₃, amino, substituted amino, cyano, or -S(O)_nR, in which R is alkyl, aryl, or heteroaryl and n is 0, 1 or 2. Heterocyclic groups can have a single ring or multiple condensed rings. Preferred heterocyclics include tetrahydrofuranyl, morpholino, piperidiny, and the like.

The term "thiol" refers to the group -SH.

The term "substituted alkylthio" refers to the group -S-substituted alkyl.

The term "heteroarylthiol" refers to the group -S-heteroaryl wherein the heteroaryl group is as defined above including optionally substituted heteroaryl groups as also defined above.

The term "sulfoxide" refers to a group -S(O)R, in which R is alkyl, aryl, or heteroaryl. "Substituted sulfoxide" refers to a group -S(O)R, in which R is substituted alkyl, substituted aryl, or substituted heteroaryl, as defined herein.

The term "sulfone" refers to a group -S(O)₂R, in which R is alkyl, aryl, or heteroaryl. "Substituted sulfone" refers to a group -S(O)₂R, in which R is substituted alkyl, substituted aryl, or substituted heteroaryl, as defined herein.

The term "keto" refers to a group -C(O)-. The term "thiocarbonyl" refers to a group -C(S)-. The term "carboxy" refers to a group -C(O)-OH.

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not.

The terms "compound of Formula I" and "compound of Formula II" are intended to encompass the compounds of the invention as disclosed, and the pharmaceutically acceptable salts, pharmaceutically acceptable esters, and prodrugs of such compounds.

The term "therapeutically effective amount" refers to that amount of an active ingredient (A₁-agonist, beta-blocker, calcium channel blocker, cardiac glycoside) that is sufficient to effect

treatment, as defined below, when administered to a mammal in need of such treatment. The therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by a prescribing physician.

- 5 The term “therapeutically effective minimal dose” or “low dose” of an A₁ adenosine receptor agonist refers to a dose level of an A₁ adenosine receptor agonist that is generally considered to be below the therapeutically effective amount as defined above, but is sufficient to provide effective treatment when administered in conjunction with a “therapeutically effective minimal dose” or “low dose” of a beta blocker, calcium channel blocker, or a cardiac glycoside.
- 10 For example, a therapeutically effective minimal dose of CVT-3619 is one that would not normally be considered to be useful in the treatment of arrhythmia, but is now found to be useful in the treatment of arrhythmia when administered in conjunction with a therapeutically effective minimal dose of a beta blocker, because of the synergistic effect obtained upon combining an A₁-agonist with a beta blocker. The therapeutically effective minimal dose will vary depending
- 15 upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by a prescribing physician.

- A therapeutically effective minimal dose of an adenosine A₁ receptor agonist is administered “in conjunction with” a therapeutically effective minimal doses of a beta blocker,
- 20 or a calcium channel blocker, or a cardiac glycoside. In this context, the word “conjunction” means that the doses may be administered together at the same time, for example in a single pill or solution, or administered separately at the same time, or administered at different times.

 The term “treatment” or “treating” means any treatment of a disease in a mammal, including:

- 25 (i) preventing the disease, that is, causing the clinical symptoms of the disease not to develop;
- (ii) inhibiting the disease, that is, arresting the development of clinical symptoms; and/or
- (iii) relieving the disease, that is, causing the regression of clinical symptoms.

- As used herein, the term “agonist” refers to the ability of a compound to interact with a
- 30 receptor and evoke a maximal effect. This effect is known as the intrinsic efficacy. In contrast, “partial agonists” interact with adenosine A₁ receptors but produce a less than maximal response.

- The term “adenosine A₁ receptor agonist” refers to an agent that binds to adenosine A₁ receptors thereby producing a negative dromotropic effect. For example, CVT-3619 is a partial A₁-adenosine receptor agonist - it has a rate dependent effect upon AV nodal conduction. It
- 35 increases AV-node refractoriness, and thus reduces ventricular rate during atrial

tachyarrhythmia. A₁ agonists also act to inhibit the release of norepinephrine from the pre-synaptic nerve terminal, and to inhibit the uptake of norepinephrine at the post-synaptic nerve terminal.

The term "beta-blocker" refers to an agent that binds to a beta-adrenergic receptor and inhibits the effects of beta-adrenergic stimulation. Beta-blockers increase AV nodal conduction. In addition, Beta-blockers decrease heart rate by blocking the effect of norepinephrine on the post synaptic nerve terminal that controls heart rate. Beta blockers also decrease intracellular Ca⁺⁺ overload, which inhibits after-depolarization mediated automaticity. Examples of beta blockers include atenolol, esmolol, sotalol, propranolol, bopindolol, carteolol, oxprenolol, penbutolol, carvedilol, medroxalol, bucindolol, levobunolol, metipranolol, betaxolol, celiprolol, and propafenone.

The term "calcium channel blocker" refers to an agent that blocks voltage-dependent "L-type calcium channel. They are used in treatment of heart diseases, including cardiac arrhythmia - they have a rate dependent effect upon AV nodal conduction. Examples of calcium channel blockers include amlodipine, bepridil, diltiazem, felodipine, isradipine, nicardipine, nifedipine, nimodipine and verapamil.

The term "cardiac glycoside" refers to a compound with a steroidal nucleus and a lactone ring, and usually has one or more sugar residues. They are used in treatment of heart diseases, including cardiac arrhythmia - they have a rate dependent effect upon AV nodal conduction. Examples of cardiac glycosides include digoxin and digitoxin.

The term "synergistic" effect means a result produced by a combination of drugs that is greater than that produced by each drug alone.

In many cases, the compounds of this invention are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto. The term "pharmaceutically acceptable salt" refers to salts that retain the biological effectiveness and properties of the compounds of Formula I, and which are not biologically or otherwise undesirable. Pharmaceutically acceptable base addition salts can be prepared from inorganic and organic bases. Salts derived from inorganic bases, include by way of example only, sodium, potassium, lithium, ammonium, calcium and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines, such as alkyl amines, dialkyl amines, trialkyl amines, substituted alkyl amines, di(substituted alkyl) amines, tri(substituted alkyl) amines, alkenyl amines, dialkenyl amines, trialkenyl amines, substituted alkenyl amines, di(substituted alkenyl) amines, tri(substituted alkenyl) amines, cycloalkyl amines, di(cycloalkyl) amines, tri(cycloalkyl) amines, substituted cycloalkyl amines, disubstituted cycloalkyl amine, trisubstituted cycloalkyl amines, cycloalkenyl amines,

di(cycloalkenyl) amines, tri(cycloalkenyl) amines, substituted cycloalkenyl amines, disubstituted cycloalkenyl amine, trisubstituted cycloalkenyl amines, aryl amines, diaryl amines, triaryl amines, heteroaryl amines, diheteroaryl amines, triheteroaryl amines, heterocyclic amines, diheterocyclic amines, triheterocyclic amines, mixed di- and tri-amines where at least two of the substituents on the amine are different and are selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic, and the like. Also included are amines where the two or three substituents, together with the amino nitrogen, form a heterocyclic or heteroaryl group.

Specific examples of suitable amines include, by way of example only, isopropylamine, trimethyl amine, diethyl amine, tri(iso-propyl) amine, tri(n-propyl) amine, ethanolamine, 2-dimethylaminoethanol, tromethamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, N-alkylglucamines, theobromine, purines, piperazine, piperidine, morpholine, N-ethylpiperidine, and the like.

Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids. Salts derived from inorganic acids include hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Salts derived from organic acids include acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluene-sulfonic acid, salicylic acid, and the like.

As used herein, "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The term "compound of Formula I" or "compound of Formula II" is intended to encompass the compounds of the invention as disclosed, and the pharmaceutically acceptable salts, pharmaceutically acceptable esters, and prodrugs of such compounds. Additionally, the compounds of the invention may possess one or more asymmetric centers, and can be produced as a racemic mixture or as individual enantiomers or diastereoisomers. The number of stereoisomers present in any given compound of the invention depends upon the number of asymmetric centers present (there are 2^n stereoisomers possible where n is the number of

asymmetric centers). The individual stereoisomers may be obtained by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolution of the compound of Formula I or Formula II by conventional means. The individual stereoisomers (including individual enantiomers and diastereoisomers) as well as racemic and non-racemic mixtures of stereoisomers are encompassed within the scope of the present invention, all of which are intended to be depicted by the structures of this specification unless otherwise specifically indicated.

"Isomers" are different compounds that have the same molecular formula.

"Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The term "(±)" is used to designate a racemic mixture where appropriate.

"Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other.

The absolute stereochemistry is specified according to the Cahn-Ingold-Prelog R-S system. When the compound is a pure enantiomer the stereochemistry at each chiral carbon may be specified by either R or S. Resolved compounds whose absolute configuration is unknown are designated (+) or (-) depending on the direction (dextro- or laevorotary) which they rotate the plane of polarized light at the wavelength of the sodium D line.

Pharmaceutical Compositions

The two components of the invention, an A₁-adenosine receptor agonist and a beta-blocker, calcium channel blocker, or a cardiac glycoside, may be administered as a pharmaceutical composition that contains a physical mixture of the two components, but is preferably administered as two separate pharmaceutical compositions. Such separate compositions are preferably administered concurrently, but may also be administered at different times. This invention therefore provides pharmaceutical compositions that contain, as the active ingredient, one or two of the components, or a pharmaceutically acceptable salt or ester thereof, and one or more pharmaceutically acceptable excipients, carriers, including inert solid diluents and fillers, diluents, including sterile aqueous solution and various organic solvents, permeation enhancers, solubilizers and adjuvants. Such compositions are prepared in a manner well known in the pharmaceutical art (see, e.g., Remington's Pharmaceutical Sciences, Mace Publishing Co., Philadelphia, PA 17th Ed. (1985) and "Modern Pharmaceutics", Marcel Dekker, Inc. 3rd Ed. (G.S. Banker & C.T. Rhodes, Eds.).

Administration

The components may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, for example as described in those patents and patent applications incorporated by reference, including rectal, buccal,
5 intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, orally, topically, as an inhalant, or via an impregnated or coated device such as a stent, for example, or an artery-inserted cylindrical polymer.

One mode for administration is parental, particularly by injection. The forms in which
10 the novel compositions of the present invention may be incorporated for administration by injection include aqueous or oil suspensions, or emulsions, with sesame oil, corn oil, cottonseed oil, or peanut oil, as well as elixirs, mannitol, dextrose, or a sterile aqueous solution, and similar pharmaceutical vehicles. Aqueous solutions in saline are also conventionally used for injection, but less preferred in the context of the present invention. Ethanol, glycerol, propylene glycol,
15 liquid polyethylene glycol, and the like (and suitable mixtures thereof), cyclodextrin derivatives, and vegetable oils may also be employed. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens,
20 chlorobutanol, phenol, sorbic acid, thimerosal, and the like.

Sterile injectable solutions are prepared by incorporating the component in the required amount in the appropriate solvent with various other ingredients as enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion
25 medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral administration is another route for administration of the components.
30 Administration may be via capsule or enteric coated tablets, or the like. In making the pharmaceutical compositions that include at least one compound of Formula I or II, the active ingredient is usually diluted by an excipient and/or enclosed within such a carrier that can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material (as above), which acts as a vehicle, carrier or
35 medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills,

powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, sterile injectable solutions, and sterile packaged powders.

5 Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; 10 preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents.

 The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art. Controlled release drug delivery systems for oral administration 15 include osmotic pump systems and dissolutional systems containing polymer-coated reservoirs or drug-polymer matrix formulations. Examples of controlled release systems are given in U.S. Patent Nos. 3,845,770; 4,326,525; 4,902,514; and 5,616,345. Another formulation for use in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the 20 compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent Nos. 5,023,252, 4,992,445 and 5,001,139. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

 The compositions are preferably formulated in a unit dosage form. The term "unit 25 dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient (e.g., a tablet, capsule, ampoule). The compounds of Formula I and II are effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It will be 30 understood, however, that the amount of the compound of Formula I actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

35 For preparing solid compositions such as tablets, the principal active ingredient is mixed

with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into
5 equally effective unit dosage forms such as tablets, pills and capsules.

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action, or to protect from the acid conditions of the stomach. For example, the tablet or pill can comprise an inner dosage and an outer dosage element, the latter being in the form of an envelope over the former. The two
10 elements can be separated by an enteric layer that serves to resist disintegration in the stomach and permit the inner element to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

15 Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described supra. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable
20 solvents may be nebulized by use of inert gases. Nebulized solutions may be inhaled directly from the nebulizing device or the nebulizing device may be attached to a face mask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices that deliver the formulation in an appropriate manner.

25 The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate
30 that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLES

The beta blockers, calcium channel blockers, and cardiac glycosides of this invention are
35 well known in the art, and are commercially available. The compounds of Formula I may be

prepared by conventional methods, in the manner disclosed in US Patent No. 5,789,416, the entire disclosure of which is hereby incorporated by reference. For example, the preferred compound CVT-510 is prepared as follows.

5

EXAMPLE 1

A. Preparation of (3-(S)-aminotetrahydrofuranyl)purine riboside

Step 1. Resolution of 3-(S)-aminotetrahydrofuran

A mixture of 3-aminotetrahydrofuran hydrochloride (0.5 GM, 4 mmol) and
10 (S)-(+)-10-camphorsulfonyl chloride (1.1 gm, 4.4 mmol) in pyridine (10 ml) was stirred for 4 hours at room temperature and then concentrated. The residue was dissolved in ethyl acetate and washed with 0.5N hydrochloric acid, followed by sodium bicarbonate and then brine. The organic layer was dried over magnesium sulfate, filtered, and solvent removed from the filtrate under reduced pressure to provide 1.17 g of a brown oil, which was chromatographed on silica
15 gel (25% to 70% ethyl acetate/hexanes). The white solid obtained was repeatedly recrystallized from acetone to yield the (S)-camphorsulfonate of 3-(S)-aminotetrahydrofuran.

Step 2. Preparation of 3-(S)-aminotetrahydrofuran hydrochloride

The (S)-camphorsulfonate of 3-(S)-aminotetrahydrofuran (170 mg, 0.56 mmol) was
20 dissolved in concentrated hydrochloric acid/acetic acid (2 mL each), and stirred for 20 hours at room temperature. The reaction mixture was washed three times with methylene chloride (10 ml), and the combined extracts concentrated to dryness under reduced pressure, to give 75 mg of 3-(S)-aminotetrahydrofuran, as a white solid.

25 Step 3. Preparation of 6-(3-(S)-aminotetrahydrofuranyl)purine riboside

A mixture of 6-chloropurine riboside (30 mg, 0.10 mmol), 3-(S)-aminotetrahydrofuran hydrochloride (19 mg, 0.15 mmol), and triethylamine (45 ml, 0.32 mmol) in methanol (0.5 ml) was heated to 80° C for 18 hours. The mixture was cooled, concentrated and chromatographed with 95/5 (methylene chloride/methanol), to give 8 mg of 6-(3-(S)-
30 aminotetrahydrofuranyl)purine riboside, as a white solid.

B. Preparation of (3-(R)-aminotetrahydrofuranyl)purine riboside (CVT-510)


Similarly, following steps 1-3 above, but replacing (S)-(+)-10-camphorsulfonyl chloride with (R)-(-)-10-camphorsulfonyl chloride, the following compound was prepared:
35 6-(3-(R)-aminotetrahydrofuranyl)purine riboside (CVT-510).

Similarly, other enantiomers of the compounds of Formula I are prepared.

The compounds of Formula II may be prepared by conventional methods, in the manner disclosed in US Patent Application Serial No. 10/194,335, the entire disclosure of which is hereby incorporated by reference. For example, the preferred compound CVT-3619 is prepared as follows.

EXAMPLE 2

10 Preparation of 6-(6-Chloropurine-9-yl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl]methanol




(2)

To a solution of 2-(6-chloropurin-9-yl)-5-hydroxymethyltetrahydrofuran-3,4-diol (a compound of formula (1)) (4.9 g, 17.1 mmol) and 2,2-dimethoxypropane (10.5mL, 84.7 mmol) in dimethylformamide (100 mL) was added p-toluenesulfonic acid (325 mg, 1.71 mmol). After stirring for 24 hours at 70°C, the reaction was concentrated in vacuo and the residue purified by flash column chromatography (70%EtOAc/Hexanes) to give 6-(6-chloropurine-9-yl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl]methanol, a compound of formula (2), as an off-white solid (2). (3.8g, 68%) ¹H NMR (CDCl₃) δ 1.4 (s, 3H), 1.65 (s, 3H), 3.8-4.0 (dd, 2H), 4.6 (s, 1H), 5.1-5.3(m, 2H), 6.0(d, 1H), 8.25(s, 1H), 8.8(s, 1H).

EXAMPLE 3

5 Preparation of 1-{[(2S,1R,4R,5R)-4-(6-chloropurin-9-yl)-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl]methylthio}-2-fluorobenzene



10 To a solution of 6-(6-chloropurin-9-yl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl]methanol, a compound of formula (2) (0.48g, 1.47 mmoles) in 20mL of tetrahydrofuran was added triphenylphosphine (0.77g, 2.94mmoles) and diethylazodicarboxylate (0.47mL, 2.94mmoles), and the mixture stirred for 5 minutes. 2-Fluorothiophenol (0.31mL, 2.94mmoles) was then added, and the mixture was stirred under reflux. After 72 hours of reflux, the reaction


15 was concentrated in vacuo and the residue purified by flash column chromatography (20%EtOAc/Hexanes) to give 1-{[(2S,1R,4R,5R)-4-(6-chloropurin-9-yl)-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl]methylthio}-2-fluorobenzene, a compound of formula (3), as a clear viscous oil (3). (0.25g, ~40%) ¹H NMR (CDCl₃) δ 1.4 (s, 3H), 1.6 (s, 3H), 3.2 (m, 2H), 4.6 (t, 1H), 5.1 (m, 1H), 5.5 (m, 1H), 6.0 (d, 1H), 7.0 (m, 2H), 7.2 (m, 1H), 7.4 (m, 1H), 8.25 (s, 1H),

20 8.75 (s, 1H).

25

EXAMPLE 4Preparation of (9-{(4S,1R,2R,5R)-4-[(2-fluorophenylthio)methyl]-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl}purin-6-yl)cyclopentylamine

To a solution of 1-{[(2S,1R,4R,5R)-4-(6-chloropurin-9-yl)-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl]methylthio}-2-fluorobenzene, a compound of formula (3), (0.125g, 2.86mmoles) in 10mL of ethanol and 1mL of triethylamine was added cyclopentylamine in excess, and the mixture refluxed under nitrogen for 24 hours. The solvent was removed under reduced pressure, and the residue was purified by preparative TLC using 1:1 EtOAc:Hexanes to give (9-{(4S,1R,2R,5R)-4-[(2-fluorophenylthio)methyl]-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl}purin-6-yl)cyclopentylamine, a compound of formula (4), as a yellow oil (80mg, 56%) ¹H NMR (CDCl₃) δ 1.4 (s, 3H), 1.6 (s, 3H), 1.6-2.4 (m, 6H), 3.15-3.25 (m, 2H), 4.1 (bs, 1H), 4.4 (t, 1H), 5.1 (m, 1H), 5.5 (m, 1H), 6.0 (d, 1H), 6.2 (bs, 1H), 7.0 (m, 2H), 7.2 (m, 1H), 7.4 (m, 1H), 7.8 (s, 1H), 8.25 (s, 1H).

EXAMPLE 5Preparation of (4S,5S,3R)-2-[6-(cyclopentylamino)purin-9-yl]-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol

Formula I

20

(9-{(4S,1R,2R,5R)-4-[(2-fluorophenylthio)methyl]-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-yl}purin-6-yl)cyclopentylamine, a compound of formula (4) (50mg) was dissolved in a mixture of acetic acid (8 mL) and water (2 mL) and heated at 90 C for 16 hours. Solvents were removed under reduced pressure, and the residue was purified by preparative TLC [methanol-dichloromethane(1:9)] to afford (4S,5S,3R)-2-[6-(cyclopentylamino)purin-9-yl]-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol, a compound of

25

Formula I, ^1H NMR (CDCl_3) δ 1.6-2.4 (m, 6H), 3.15-3.25 (m, 2H), 4.1 (bs, 1H), 4.4-4.65 (m, 4H), 6.0 (d, 1H), 6.8 (bs, 1H), 7.05 (m, 2H), 7.2 (m, 1H), 7.4 (m, 1H), 7.8 (s, 1H), 8.25 (s, 1H).

5

EXAMPLE 6

Studies in awake rats:

Methods

Rats (Sprague Dawley) weighing 300-400 gms were purchased from Simonsen laboratories. CVT-3619 was dissolved in DMSO and further diluted in saline. CVT-510 was dissolved in saline. Ketamine was purchased from Fort Dodge Animal Health, Xylazine from Bayer and Acepromazine Maleate from Fermenta Animal Health Co. Metoprolol and propranolol were purchased from SIGMA. Esmolol was obtained from a local pharmacy.

Telemetry Studies

For these studies, rats were instrumented with radiotelemetered transmitters (Data Sciences) at least 3 weeks prior to experimentation. Animals were anesthetized by peritoneal injection of a "cocktail" (1 ml/kg) containing Ketamine (75mg/ml), Xylazine (5mg/ml), and Acepromazine (1mg/ml). After 20-30 minutes of induction of anesthesia, a midline laparotomy was performed. The transmitter for recordings of ECG, blood pressure and body temperature was placed in the abdominal cavity, and secured to the abdominal muscles. Two electrocardiographic leads were tunneled through the subcutaneous - one toward the upper left shoulder and the other to the right thigh, and secured with sutures. A fluid filled sensor catheter was inserted in the descending aorta above the iliac bifurcation for measurement of blood pressure. The tip of the telemetry catheter was located in the abdominal aorta just caudal to the renal arteries. Once the transmitter and leads were in place and determined to be functioning properly, the abdominal wall was sutured. After recovery from anesthesia, the rats were housed individually in cages placed on their respective receivers. The ECG, blood pressure and temperature were recorded and heart rate measured by a Dataquest ART Gold system (Version 2.2; Data Sciences Intl). The system consisted of a transmitter, i.e., biopotential sensor (Model TL11M2-C50-PXT), receivers (Model RPC-1), a consolidation matrix (BCM 100), a personal computer (Compaq DeskPro Series 3574) and Dataquest 4 software. Heart rate, blood pressure and temperature were measured at 5-minute intervals. Each recording lasted 10 seconds and all cardiac cycles within this period were averaged. Animals were given various drugs in a randomized manner after recording the baseline data for at least two hours.

Implantation of Osmotic Pump

A subgroup of animals was implanted with osmotic pumps containing CVT-510 for combination experiments. After 7 days or more of having been implanted with radio-telemetry transmitters for recording ECG (L-2), and verifying that the transmitters were functioning, each rat was implanted with an Alzet mini-osmotic pump. Under anesthesia (see above) and sterile conditions, one Alzet mini-osmotic pump, was implanted subcutaneously (SC) in the interscapular area of each rat. The osmotic pumps were filled with either vehicle or CVT-510 (to deliver a dose of 20 µg/kg/hr).

10

Surgical insertion of catheters for determining plasma concentrations

Carotid artery was catheterized to obtain serial blood samples for the analysis of CVT-510 plasma concentrations. Animals were anesthetized by peritoneal injection of a “cocktail” (1 ml/kg) containing Ketamine (75mg/ml), Xylazine (5mg/ml), and Acepromazine (1mg/ml). After a 20-30 min. of induction of anesthesia, a midline incision was made in the neck region to expose the external carotid artery. A tunnel is made for the catheter using blunt dissection in the subcutaneous pocket on the dorsal section of the neck where it is externalized. The carotid artery was cannulated with 24-gauge catheters sampling of blood for determination of plasma levels of CVT-510. Externally, the catheter is tied at the back of the neck and a piece of suture is tied around the knot leaving both ends about 2 inches long for retrieval from under the skin. The knotted catheter is retracted back under the skin to prevent being pulled out by the rat. The incision is then cleaned with saline, closed with wound clips, and an antibiotic (0.4 ml of a 40mg/ml solution of gentamicin) is given I.V. Animals were allowed to recover for at least 48 hrs before performing the experiment. On the day of the experiment, an injection plug was attached to a 19-gauge IV set, filled with 0.1% heparinized saline and the needle end was inserted into the catheters. Animals were given either a saline or metoprolol injection 1 hr prior to CVT-510 injection. About 400 µl of blood was drawn from the line in the carotid artery and 400 ul saline flushed in to replace blood volume at predetermined time points. Plasma was separated and saved at -80°C for analysis of CVT-510 levels.

20
25
30

Determination of plasma concentrations of CVT-510

CVT-510 plasma level analysis was performed as follows. Briefly, 50 µL of plasma sample was precipitated with 400 µL of acetonitrile:methanol (90:10) containing internal standard. The filtrates were evaporated to dryness and reconstituted in 100 µL of 90:10 water:methanol. Concentration of CVT-510 in protein precipitation filtrates were analyzed by

35

LC-MS-MS using a Waters Alliance 2690 HPLC system (Millford, MA) coupled to a Waters/Micromass Quattro Ultima triple quadrupole mass spectrometer (Millford, MA). Calibration curves were constructed by plotting peak area ratios of the analyte to internal standard against concentration, using a weighted (1/X) linear regression model in all instances.

5

Data Analysis

The slowing of heart rate caused by each treatment was quantified by determining the area under the curve (AUC), using the trapezoidal method for calculations. Data used for analysis was the area under/over the curve calculated using change from baseline data (untreated animals). The data was analyzed for the magnitude as well as the duration of bradycardia caused by each treatment. The AUC values for various doses for each monotherapy were compared using one-way ANOVA followed by Tukey's test for multiple comparisons. The significance level was set at $p < 0.05$ for all comparisons.

15

The results obtained from the above studies are shown in Figure form, as follows.

Figure 1 shows the effect of CVT-3619 (a partial A_1 receptor agonist) on heart rate with and without propranolol in awake rats instrumented with telemetry radiotransmitters. CVT-3619 at dose of (0.5 mg/kg, ip) by itself had minimal effect on heart rate. However when given in the presence of a beta blocker (propranolol, 10 mg/kg, ip), there is significant lowering of heart rate below baseline as compared to CVT-3619 alone. Propranolol was given 10 minutes prior to CVT-3619 injection.

Figure 2 shows the summary of data obtained from telemeterized awake rats. The data was quantified as area under the curve (AUC) and presented as a decrease in total number of heart beats for a 2 hour period of time caused by CVT-3619 (given ip) alone at two different doses and in the presence of propranolol (10 mg/kg, ip given 10 minutes prior to CVT-3619). The combination of CVT-3619 and propranolol results in a synergistic effect on heart rate, as the effect observed with combination is much greater than the calculated sum of the effect of each agent.

30

Figure 3 shows the effect of CVT-510 and esmolol alone and given together as a mixture via ip injection to awake rats instrumented with telemetry radiotransmitters. CVT-510 at 20 μ g/kg dose transiently lowers the heart rate below baseline levels whereas esmolol (10 mg/kg ip) only slightly reduces the increase in heart rate (which is caused by handling the animal). When the two agents are given in combination, the effect on heart rate is much greater. The

35

combination not only increases the magnitude of the response but also increases the duration of bradycardia significantly.

5 Figure 4 shows the effect of CVT-510 alone and the effect when combined with the beta-blocker, Metoprolol, by observing the effect on heart rate in awake rats instrumented with telemetry radiotransmitters. CVT-510 (20 μ g/kg, ip) slowed the heart rate in a dose dependent manner. In the presence of Metoprolol (3 mg/kg, ip), there was a greater lowering of heart rate.

10 Figure 5. The duration of bradycardia caused by CVT-510 was analyzed by observing the time for the heart rate to return to 90% of pretreatment levels. In addition to the increase in magnitude of bradycardia, the combination of CVT-510 and metoprolol resulted in a significant increase in the duration of bradycardia as compared to that caused by CVT-510 alone.

15 Figure 6: Plasma levels of CVT-510 were determined in the absence and presence of metoprolol in a separate group of animals (figure 6). Plasma levels of CVT-510 were found to be similar in both groups even though the slowing of heart rate caused by CVT-510 was greater in the presence of metoprolol indicating that the metabolism of CVT-510 was not changed in the presence of metoprolol.

20 Figure 7: To further investigate the mechanism of interaction of CVT-510 and beta-blockers, another series of experiments were performed in which the dose of CVT-510 was kept constant while the dose of metoprolol was varied. First a full dose response curve for metoprolol (0.1-10 mg/kg, ip) alone was obtained (figure 7, ■ symbols). In the second phase of the study, animals were implanted with osmotic pumps subcutaneously containing CVT-510. CVT-510 was
25 delivered at a rate of 20 μ g/kg/hr, which yielded plasma concentration of 7.5 ± 1 ng/ml. Metoprolol dose response curve was repeated in these animals. In the presence of CVT-510, metoprolol dose response curve was shifted to the left and downward (figure 7, ● symbols).

30 Figure 8: This figure represents the data shown in figure 7. The slowing of heart rate caused by CVT-510 and metoprolol was quantified by determining the area under the curve (AUC) for a period of 60 min for each treatment. The combined effect of CVT-510 and metoprolol on heart rate was found to be synergistic.

Figure 9-10: Studies in anesthetized guinea pigs.

Male guinea pigs weighing 400-450 gm were obtained from Simonsen labs and housed in the institutional laboratory animal facility. Animals were anesthetized with isoflurane in the anesthetizing chamber. After determining (by toe pinch) that the animal is adequately anesthetized, the animals was intubated with an endotracheal tube and ventilated with isoflurane and oxygen mixture using anesthesia workstation. Using sterile equipment and aseptic technique, the right carotid artery was exposed and a catheter inserted for recording of blood pressure (BP). A quadripolar electrode catheter was introduced via the right jugular vein and positioned in the right atrium and ventricle for atrial and ventricular pacing. The hearts were paced at a constant rate (330-360 bpm) to eliminate the effect of heart rate variability between animals. Another catheter was inserted into the left jugular vein and positioned in the right atrium for the administration of drugs and saline. Subcutaneous needles were used as standard electrocardiographic leads to record the electrocardiogram (ECG). After completion of surgery and instrumentation, a 20 min equilibration period was allowed before beginning the experimental protocol. The data was recorded using Power Lab data acquisition system.

Figure 9: CVT-510 (0.5 $\mu\text{g/kg}$) and metoprolol (0.1 mg/kg), which were given as an iv bolus, demonstrated an increased PR interval by 5 msec each in anesthetized guinea pigs. When the two agents were given in combination the PR interval was increased up to 15 msec. This is a synergistic effect, as the observed effect is more than the algebraic sum of the effect of each agent.

Figure 10: Similar results were obtained when CVT-510 was given in the presence of with another beta-blocker, esmolol. Esmolol was administered at three different infusion rates. CVT-510 was given 15 minutes after starting the infusion of Esmolol. Effect of CVT-510 on PR interval was increased with increasing doses of esmolol. The duration of effect was significantly prolonged in a dose-dependent manner.

Thus, it has been demonstrated that the combination of a beta-blocker and A_1 agonist results in synergistic effects on heart rate in rats and PR interval in guinea pigs. The combined effect is dependent on the dose of either agent. That is, one can achieve similar responses by keeping one agent constant and varying the other. Various routes of administration of the drugs have been tried, and different combinations. For example, one drug has been administered 10 minutes after administration of the first, and 1 hour after administration of the first. The drugs have been given as a mixture, or given separately at the same time. The response varies in magnitude, but the overall effect is same. It has also been demonstrated that the combination is

effective using both full A₁ adenosine receptor agonists and partial A₁ adenosine receptor agonists.

The combination has been shown to be effective in two different models.

- 1) Measurement of heart rate in awake rats, which is not a target for the A₁ agonists, but is used
5 as a surrogate for the effect of A₁ agonists. The advantage of this model is that the high
sympathetic tone seen in many forms of arrhythmias is simulated.
- 2) In anesthetized guinea pigs the AV nodal conduction method is useful for measuring the PR
interval, which is the target of A₁ agonists. However, the sympathetic tone is blunted in this
model due to anesthesia.

10

CVT-510 (SH)

Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	DIGOXIN+CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
digoxin#1	50	70		100	100	160	50	90	80	90		140
digoxin#2	30	40	50	60	110		50	60	70	80		100
digoxin#3	40	50	50	70	100	140	50	90	100	100	100	160
digoxin#4	60	90	120	120	120	140	60	90	120	140	130	150
digoxin#5	40	60	70	100	150	100	60	120	120	120	100	90
n	5	5	4	5	4	5	5	5	5	5	3	5
MEAN	44	62	73	90	118	130	54	90	98	106	110	128
SD	11	19	33	24	24	24	5	21	23	24	17	31
SEM	5	9	17	11	12	11	2	9	10	11	10	14

Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	ESMOLOL+CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
esmolol#1	40	60	60	100	100	110	60	80	90	100	100	100
esmolol#2	30	40	60	60	80	100	50	60	60	90	90	140
esmolol#3	50	80	90	120	140	140	50	80	130	120	140	
esmolol#4	30	40	50	70	90	90	40	50	50	70	80	80
esmolol#5	40	90	90	100	110	100	50	90	120	100	130	130
n	5	5	5	5	5	5	5	5	5	5	5	4
MEAN	38	62	70	90	104	108	50	72	90	96	108	113
SD	8	23	19	24	23	19	7	16	35	18	26	28
SEM	4	10	8	11	10	9	3	7	16	8	12	14

Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	IBUTILIDE+CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
ibutilide#1	35	40	45	60	80	140	40	80	90	90	80	
ibutilide#2	70	90	150	150	150	160	75	110	110	130	120	110
ibutilide#3	40	40	50	55	70	75	50	60	70	70	70	70
ibutilide#4	40	60	70	90	100	80	50	70	80	90	90	120
ibutilide#5	40	45	50	55	80	90	50	60	65	70	90	80
n	5	5	5	5	5	5	5	5	5	5	5	4
MEAN	45	55	73	82	96	109	53	76	83	90	90	95
SD	14	21	44	41	32	38	13	21	18	24	19	24

SEM	6	9	20	18	14	17	6	9	8	11	8	12
-----	---	---	----	----	----	----	---	---	---	----	---	----

Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	QUINIDINE + CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
quinidine#1	40	50	55	60	90	90	50	70	70	70	105	125
quinidine#2	30	40	45	45	100	100	50	70	70	65	90	130
quinidine#3	40	60	65	70	110	110	70	80	85	95	115	120
quinidine#4	40	50	90	110	130	120	60	75	115	115	120	95
quinidine#5	50	50	60	70	80	70	40	60	95	100	90	105
N	5	5	5	5	5	5	5	5	5	5	5	5
MEAN	40	50	63	71	102	98	54	71	87	89	104	115
SD	7	7	17	24	19	19	11	7	19	21	14	15
SEM	3	3	8	11	9	9	5	3	8	9	6	7

Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	PROCAINAMIDE + CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
procainamide#1	45	50	60	65	95	85	65	65	70	80	90	110
procainamide#2	50	55	70	95	105	105	55	80	110	120	120	120
procainamide#3	40	50	80	95	95	110	55	65	105	110	110	110
procainamide#4	40	50	55	65	75	75	65	50	80	90	90	90
procainamide#5	45	55	65	80	90	100	50	80	85	85	85	85
n	5	5	5	5	5	5	5	5	5	5	5	5
MEAN	44	52	66	80	92	95	58	68	90	97	99	103
SD	4	3	10	15	11	15	7	13	17	17	15	15
SEM	2	1	4	7	5	7	3	6	8	8	7	7


Animal #	baseline	CVT-510 Alone (µg/kg)					baseline	ATENOLOL + CVT-510 (µg/kg)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
atolol#2	50	70	100	180	120	120	70	120	170	180	180	220
atolol#3	45	80	100	120	120	120	60	130	150	180	180	180
atolol#4	45	50	50	75	100	85	45	60	90	110	125	130
atolol#5	45	50	70	90	130	165	50	120	160	130	160	160
atolol#6	40	50	110	115	120	120	50	65	120	120	120	210
n	5	5	5	5	5	5	5	5	5	5	5	5
MEAN	45	60	86	116	118	122	55	99	138	144	153	180

SD	4	14	25	40	11	28	10	34	33	34	29	37
SEM	2	6	11	18	5	13	4	15	15	15	13	16

Animal #	baseline	CVT-510 Alone ($\mu\text{g/kg}$)					baseline	SOTALOL+CVT-510 ($\mu\text{g/kg}$)				
		1.69	2.53	3.37	8.43	16.85		1.69	2.53	3.37	8.43	16.85
sotalol#1	40	70	110	140	140	140	60	90	140	140	140	180
sotalol#2	40	55	100	110	130	140	60	130	140	140	140	140
sotalol#3	55	60	70	80	85	130	65	85	115	110	130	190
sotalol#4	40	45	50	50	70	105	60	90	85	100	100	105
sotalol#5	40	50	110	115	120	120	70	65	120	120	120	210
n	5	5	5	5	5	5	5	5	5	5	5	5
MEAN	43	56	88	99	109	127	63	92	120	122	126	165
SD	-7	10	27	35	30	15	4	24	23	18	17	42
SEM	3	4	12	16	13	7	2	11	10	8	7	19

What is claimed is:

1. A method of treating arrhythmias in a mammal, comprising administration of a therapeutically effective minimal dose of an A₁ adenosine receptor agonist in conjunction with a therapeutically effective minimal dose of a beta blocker, calcium channel blocker, or cardiac glycoside, to a mammal in need thereof.
2. The method of claim 1, wherein the A₁ adenosine receptor agonist is a compound of compound of Formula I:



Formula I

wherein R₁ is an optionally substituted heterocyclic group.


3. The method of claim 2, wherein the compound of Formula I is administered in conjunction with a therapeutically effective minimal dose of a beta blocker.
4. The method of claim 3, wherein in Formula I R¹ is 3-tetrahydrofuranyl, 3-tetrahydrothiofuranyl, 4-pyranyl, or 4-thiopyranyl.
5. The method of claim 4, wherein the beta blocker is atenolol, esmolol, sotalol, or propranolol.
6. The method of claim 5, wherein the compound of Formula I R¹ is 6-(3-(R)-N-aminotetrahydrofuranyl)purine riboside, namely CVT-510.
7. The method of claim 6, wherein CVT-510 is present in an amount from about 0.0001-0.05 mg/kg.

8. The method of claim 6, wherein the beta blocker is present in an amount from about 0.01 to 200 mg/kg.

5 9. The method of claim 8, wherein the beta blocker is esmolol.

10. The method of claim 9, wherein CVT-510 is present in an amount from about 0.0005-0.020 mg/kg and esmolol is present in an amount from about 0.1 to 10 mg/kg.

10 11. The method of claim 1, wherein the A₁ adenosine receptor agonist is a compound of Formula II:



Formula II

wherein:

R¹ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl, or
15 optionally substituted heteroaryl;

R² is hydrogen, halo, trifluoromethyl, optionally substituted acyl, or cyano;

R³ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl;
optionally substituted heteroaryl, or optionally substituted heterocyclyl,

R⁴ and R⁵ are independently hydrogen or optionally substituted acyl; and

20 X and Y are independently a covalent bond or optionally substituted alkylene.

12. The method of claim 11, wherein the compound of Formula II is administered in conjunction with a therapeutically effective minimal dose of a beta blocker.

25 13. The method of claim 12, wherein the beta blocker is atenolol, esmolol, sotalol, or propranolol.

14. The method of claim 13, wherein R^1 is (R)-2-hydroxycyclopentyl, X and Y are covalent bonds, R^2 , R^3 , and R^4 are hydrogen, and R^5 is 2-fluorophenyl, namely 2-{6-[(1R,2R)-2-hydroxycyclopentyl]amino]purin-9-yl}(4S,5S,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol (CVT-3619).

15. The method of claim 14, wherein CVT-3619 is present in an amount from about 0.1 to 200 mg/kg.


16. The method of claim 14, wherein the beta blocker is present in an amount from about 0.01 to 100 mg/kg.

17. The method of claim 16, wherein the beta blocker is esmolol.

18. The method of claim 17, wherein CVT-3619 is present in an amount from about 0.5 to 50 mg/kg and esmolol is present in an amount from about 0.1 to 10 mg/kg.

19. A pharmaceutical composition comprising a therapeutically effective minimal dose of an A_1 adenosine receptor agonist and a therapeutically effective minimal dose of a beta blocker, and at least one pharmaceutically acceptable excipient.


20. The pharmaceutical composition of claim 19, wherein the A_1 adenosine receptor agonist is a compound of compound of Formula I:



Formula I

wherein R_1 is an optionally substituted heterocyclic group.

21. The pharmaceutical composition of claim 20, wherein the compound of Formula I is administered in conjunction with a therapeutically effective minimal dose of a beta blocker.
22. The pharmaceutical composition of claim 21, wherein in Formula I R^1 is 3-
5 tetrahydrofuranyl, 3-tetrahydrothiofuranyl, 4-pyranyl, or 4-thiopyranyl.
23. The pharmaceutical composition of claim 22, wherein the beta blocker is atenolol, esmolol, sotalol, or propranolol.
- 10 24. The pharmaceutical composition of claim 23, wherein the compound of Formula I R^1 is 6-(3-(R)-N-aminotetrahydrofuranyl)purine riboside, namely CVT-510.
25. The pharmaceutical composition of claim 24, wherein CVT-510 is present in an amount from about 0.0001-0.05 mg/kg.
- 15 26. The pharmaceutical composition of claim 24, wherein the beta blocker is present in an amount from about 0.01 to 100 mg/kg.
27. The pharmaceutical composition of claim 26, wherein the beta blocker is esmolol.
- 20 28. The pharmaceutical composition of claim 27, wherein CVT-510 is present in an amount from about 0.0005-0.02 mg/kg and esmolol is present in an amount from about 0.1 to 10 mg/kg.
29. The pharmaceutical composition of claim 19, wherein the A_1 adenosine receptor agonist
25 is a compound of Formula II:



Formula II

wherein:

R¹ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

5 R² is hydrogen, halo, trifluoromethyl, optionally substituted acyl, or cyano;

R³ is optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl; optionally substituted heteroaryl, or optionally substituted heterocyclyl,

R⁴ and R⁵ are independently hydrogen or optionally substituted acyl; and

X and Y are independently a covalent bond or optionally substituted alkylene.

10

30. The pharmaceutical composition of claim 29, wherein the compound of Formula II is administered in conjunction with a therapeutically effective minimal dose of a beta blocker.

31. The pharmaceutical composition of claim 30, wherein the beta blocker is atenolol,
15 esmolol, sotalol, or propranolol.

32. The pharmaceutical composition of claim 31, wherein R¹ is (R)-2-hydroxycyclopentyl, X and Y are covalent bonds, R², R³, and R⁴ are hydrogen, and R⁵ is 2-fluorophenyl, namely 2-{6-
20 [((1R,2R)-2-hydroxycyclopentyl)amino]purin-9-yl}(4S,5S,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol (CVT-3619).

33. The pharmaceutical composition of claim 32, wherein CVT-3619 is present in an amount from about 0.1 to 200 mg/kg.

25 34. The pharmaceutical composition of claim 32, wherein the beta blocker is present in an amount from about 0.01 to 100 mg/kg.

35. The pharmaceutical composition of claim 34, wherein the beta blocker is esmolol.
36. The pharmaceutical composition of claim 35, wherein CVT-3619 is present in an amount
5 from about 0.5 to 25 mg/kg and esmolol is present in an amount from about 0.1 to 10 mg/kg.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9





Figure 10



INTERNATIONAL SEARCH REPORT

International Classification No

PCT/US 03/12043

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K31/7076 A61K31/138 A61K31/44 A61P9/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2001/008883 A1 (BELARDINELLI LUIZ ET AL) 19 July 2001 (2001-07-19) page 5, left-hand column abstract page 1, column 2	1
Y	US 5 789 416 A (LUM ROBERT T ET AL) 4 August 1998 (1998-08-04) cited in the application page 1, left-hand column, line 55 -right-hand column, line 67 --- -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

20 August 2003

Date of mailing of the international search report

28/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Young, A

INTERNATIONAL SEARCH REPORT

Internationl Application No

PCT/US 03/12043

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SNOWDY S ET AL: "A COMPARISON OF AN A1 ADENOSINE RECEPTOR AGONIST (CVT-510) WITH DILTIAZEM FOR SLOWING OF AV NODAL CONDUCTION IN GUINEA-PIG" BRITISH JOURNAL OF PHARMACOLOGY, BASINGSTOKE, HANTS, GB, vol. 126, January 1999 (1999-01), pages 137-146, XP000992136 ISSN: 0007-1188 abstract	1
A	WO 01 23399 A (MASAMUNE HIROKO ;PFIZER PROD INC (US); DENINNO MICHAEL PAUL (US);) 5 April 2001 (2001-04-05) page 25, line 27 -page 27, line 1 claims 39,40	1-36

INTERNATIONAL SEARCH REPORT

Interr application No.
PCT/US 03/12043

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-18 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

 Internat
 PCT/US 03/12043

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2001008883 A1	19-07-2001	US 6294522 B1	25-09-2001
		US 6258793 B1	10-07-2001
		AU 4138601 A	12-06-2001
		AU 4510101 A	18-06-2001
		CA 2394861 A1	14-06-2001
		CN 1402732 T	12-03-2003
		EP 1237898 A2	11-09-2002
		JP 2003516413 T	13-05-2003
		NO 20022573 A	30-05-2002
		TR 200201470 T2	23-12-2002
		WO 0142265 A2	14-06-2001
		WO 0140799 A2	07-06-2001
		US 2002045595 A1	18-04-2002
		AU 1937201 A	12-06-2001
		AU 2055201 A	12-06-2001
		WO 0140244 A1	07-06-2001
		WO 0140245 A1	07-06-2001
		US 2002037872 A1	28-03-2002
		AU 1811701 A	12-06-2001
		AU 4138701 A	12-06-2001
		AU 761029 B2	29-05-2003
		AU 7938301 A	21-03-2002
		BR 0016126 A	06-08-2002
		CA 2389264 A1	07-06-2001
		CN 1378550 T	06-11-2002
		CZ 20013705 A3	17-04-2002
		EP 1233973 A1	28-08-2002
		JP 2003516325 T	13-05-2003
		NO 20015373 A	29-11-2001
		TR 200103129 T2	21-10-2002
		WO 0140246 A1	07-06-2001
		WO 0140243 A2	07-06-2001
US 5789416 A	04-08-1998	AT 202361 T	15-07-2001
		AT 234323 T	15-03-2003
		AU 726597 B2	16-11-2000
		AU 4080997 A	19-03-1998
		BR 9711444 A	18-01-2000
		CN 1234803 A	10-11-1999
		CZ 9900614 A3	13-10-1999
		DE 69705312 D1	26-07-2001
		DE 69705312 T2	11-10-2001
		DE 69719816 D1	17-04-2003
		DK 920438 T3	10-09-2001
		DK 992510 T3	07-07-2003
		EP 1081155 A2	07-03-2001
		EP 0920438 A2	09-06-1999
		EP 0992510 A1	12-04-2000
		ES 2157593 T3	16-08-2001
		ES 2189330 T3	01-07-2003
		GR 3036332 T3	30-11-2001
		HU 0001640 A2	28-05-2001
		JP 2000501426 T	08-02-2000
		JP 3157842 B2	16-04-2001
		KR 2000035867 A	26-06-2000
		NO 990787 A	19-02-1999
		NO 20020758 A	19-02-1999
		NO 20020759 A	19-02-1999

INTERNATIONAL SEARCH REPORT

International	Application No
PCT/US 03/12043	

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5789416	A		NZ 334095 A	27-10-2000
			PL 331853 A1	16-08-1999
			PT 920438 T	30-10-2001
			TR 9900377 T2	21-05-1999
			TR 200202287 T2	21-01-2003
			WO 9808855 A2	05-03-1998
<hr/>				
WO 0123399	A	05-04-2001	AU 7035200 A	30-04-2001
			BG 106636 A	31-01-2003
			BR 0014384 A	02-07-2002
			CA 2386079 A1	05-04-2001
			CN 1374967 T	16-10-2002
			CZ 20021020 A3	16-04-2003
			EE 200200172 A	16-06-2003
			EP 1216257 A1	26-06-2002
			HU 0202807 A2	28-12-2002
			WO 0123399 A1	05-04-2001
			JP 2003510331 T	18-03-2003
			NO 20021474 A	22-05-2002
			TR 200200843 T2	22-07-2002
<hr/>				